

MYERS 1086-1087

Terms & Conditions


Electronic Supporting Information files are available without a subscription to ACS Web Editions. The American Chemical Society holds a copyright ownership interest in any copyrightable Supporting Information. Files available from the ACS website may be downloaded for personal use only. Users are not otherwise permitted to reproduce, republish, redistribute, or sell any Supporting Information from the ACS website, either in whole or in part, in either machine-readable form or any other form without permission from the American Chemical Society. For permission to reproduce, republish and redistribute this material, requesters must process their own requests via the RightsLink permission system. Information about how to use the RightsLink permission system can be found at <http://pubs.acs.org/page/copyright/permissions.html>.



ACS Publications

MOST TRUSTED. MOST CITED. MOST READ.

J-1687-m1





^1H NMR (400 MHz, C_6D_6):
5.68 (dd, 1H, $J \sim 1.5, 2.0$ Hz, H10)
5.20 (bs, 1H, H8)
4.42 (d, 1H, $J = 12.4$ Hz, H13)
3.81 (d, 1H, $J = 12.4$ Hz, H13)
3.09 (s, 1H, H5)
2.24-2.31 (m, 1H, H11)
1.93-2.03 (m, 3H, H11, H12)
1.11 (s, 9H, $\text{O}_2\text{Ct-Bu}$)
0.26 (s, 9H, $\text{Si}(\text{CH}_3)_3$)

FTIR (neat film, cm^{-1}): 3444(b), 2210(w), 1738(s), 1252(m), 1142(s).

TLC (1:1 Et_2O :pet ether): $R_f = 0.32$


HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 389.1784 $[\text{MH}]^+$
found: 389.1795



J-1087-m2

J-1087-m3



¹H NMR (400 MHz, C₆D₆):




- 5.92 (bd, 1H, *J*=4.4 Hz, H10)
- 5.02 (bs, 1H, H8)
- 4.19 (d, 1H, *J*=12.2 Hz, H13)
- 3.93 (d, 1H, *J*=12.2 Hz, H13)
- 3.18 (d, 1H, *J*=1.5 Hz, H5)
- 1.95-2.03 (m, 1H, H11)
- 1.62-1.75 (m, 2H, H12)
- 1.15-1.24 (m, 1H, H11)
- 1.09 (s, 9H, O₂Ct-Bu)
- 0.15 (s, 9H, Si(CH₃)₃)

FTIR (neat film, cm⁻¹): 1784(s), 1741(m), 1146(s)

TLC (20% EtOAc-Hexanes): *R*_f = 0.53


HR FAB MS,

(matrix: Nitrobenzyl alcohol): calcd: 485.1607 [MH]⁺
found: 485.1617



744-6801-f

J-1087-m5






^1H NMR (400 MHz, C_6D_6): 7.78 (ddd, 2H, $J=1.5, 8.3, 9.8$ Hz, *o*-Ar)
6.91-7.06 (m, 3H, Ar)
6.16-6.20 (m, 1H, H8)
4.37-4.45 (m, 1H, H10)
4.19 (d, 1H, $J=12.4$ Hz, H13)
3.83 (d, 1H, $J=12.4$ Hz, H13)
3.16 (d, 1H, $J=0.7$ Hz, H5)
1.68-1.74 (m, 2H, H11 or H12)
1.39-1.46 (m, 2H, H11 or H12)
1.11 (s, 9H, $\text{O}_2\text{Ct-Bu}$)

FTIR (neat film, cm^{-1}): 2190(w), 1733(s), 1354(m), 1277(s), 1132(s)

TLC (1:2 EtOAc:Hexanes): $R_f = 0.23$


HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 439.1311 $[\text{M} + \text{O} + \text{H}]^+$
found: 439.1335



964-6801-7

J-1087-m7



^1H NMR (400 MHz, C_6D_6):




- 5.03 (bdd, 1H, $J=1.7, 7.6$ Hz, H8)
- 4.58-4.64 (m, 1H, H10)
- 4.28 (d, 1H, $J=12.4$ Hz, H13)
- 3.94 (d, 1H, $J=12.4$ Hz, H13)
- 3.15 (d, 1H, $J=1.7$ Hz, H5)
- 2.18-2.27 (m, 1H, H12)
- 1.95-2.03 (m, 1H, H12)
- 1.73-1.82 (m, 1H, H11)
- 1.40-1.50 (m, 1H, H11)
- 1.10 (s, 9H, $\text{O}_2\text{Ct-Bu}$)
- 1.04 (d, 1H, $J=9.3$ Hz, OH)
- 0.90 (s, 9H, $\text{Si}(\text{CH}_3)_3$)
- 0.03 (s, 3H, SiCH_3)
- 0.01 (s, 3H, SiCH_3)

FTIR (neat film, cm^{-1}): 3422(b), 2188(w), 1738(s), 1149(s)

TLC (20% EtOAc-Hexanes): $R_f = 0.37$


HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 431.2254 $[\text{MH}]^+$
found: 431.2240



844-6801-f

J-1087-m9



^1H NMR (400 MHz, C_6D_6):




- 5.78 (bs, 1H, H12)
- 5.49 (bs, 1H, H8)
- 4.37 (d, 1H, $J=12.2$ Hz, H13)
- 4.25 (m, 1H, H10)
- 3.96 (d, 1H, $J=12.2$ Hz, H13)
- 3.46 (s, 1H, H5)
- 1.93-2.06 (m, 2H, H11)
- 1.12 (s, 9H, $\text{O}_2\text{Ct-Bu}$)
- 0.85 (s, 9H, $\text{Si}(\text{CH}_3)_3$)
- 0.11 (s, 3H, SiCH_3)
- 0.13 (s, 3H, SiCH_3)

FTIR (neat film, cm^{-1}): 2177(w), 1733(s)



TLC (20% EtOAc-Hexanes): $R_f = 0.46$

HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 413.2148 $[\text{MH}]^+$
found: 413.2147



J-1087-m10



Synthetic (7)
¹³C NMR (300 MHz, C₆D₆)



176.6 (C15)
166.0 (C9)
141.9 (C12)
124.5 (C1)
99.8 (C8) [†]
98.0 (C6) [†]
96.4 (C3) [†]
90.2 (C7) [†]
84.7 (C2) [†]
72.6 (C10)
62.5 (C13) [†]
61.3 (C4) [†]
53.9 (C5)
40.7 (CO-C-(CH ₃) ₃) [†]
38.5 (C11) [†]
26.8 (CO-C-(CH ₃) ₃) [†]
25.5 (Si-C-(CH ₃) ₃) [†]
17.7 (Si-C-(CH ₃) ₃)
- 5.0 (SiCH ₃)
- 5.3 (SiCH ₃)

NCS Chromophore (1)
¹³C NMR (400 MHz,
 1:1 CD₃CO₂D:CD₃OD)[‡]

-
160.2 (C9)
139.4 (C12)
129.8 (C1)
106.5 (C8)
99.7 (C6)
97.6 (C3)
90.7 (C7)
87.5 (C2)
82.2 (C10)
76.1 (C13)
63.8 (C4)
55.2 (C5)
-
81.6 (C11)
-
-
-
-
-
-


[†]tentative assignments

[‡]Edo, K.; Mizugaki, M.; Koide, Y.; Seto, H.; Furihata, K.; Otake, N.; Ishida, N.
Tetrahedron Lett. **1985**, 26, 331.



✓-1087-m12

J-1087-m13



^1H NMR (400 MHz, CDCl_3):




- 7.40 (s, 1H, H2)
- 7.12 (s, 1H, H8)
- 6.71 (d, 1H, $J=5.9$ Hz, H6)
- 6.34 (d, 1H, $J=5.9$ Hz, H5)
- 5.40 (t, 1H, $J=6.8\pm 0.3$ Hz, H10)
- 4.52 (dd, 1H, $J=1.7, 6.1$ Hz, H12)
- 4.33 (d, 1H, $J=11.4$ Hz, H13)
- 4.10 (d, 1H, $J=11.4$ Hz, H13)
- 3.72 (s, 3H, CO_2CH_3)
- 3.22 (AB, 2H, $J=14.9$ Hz, SCH_2)
- 2.50 (ddd, 1H, $J=1.7, 6.4, 8.4$ Hz, H11)
- 2.32-2.42 (m, 1H, H11)
- 1.22 (s, 9H, $\text{O}_2\text{Ct-Bu}$)
- 0.92 (s, 9H, $\text{Si}(\text{CH}_3)_3$)
- 0.17 (s, 3H, SiCH_3)
- 0.14 (s, 3H, SiCH_3)

FTIR (neat film, cm^{-1}): 3452(b), 1734(s), 1717(s), 1532(m), 1349(m)

TLC (1:2 EtOAc:Hexanes): $R_f = 0.37$


HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 519.2237 $[\text{M} - \text{H}]^+$
found: 519.2271



7100-6801-5

J-1087-m15



^1H NMR (400 MHz, CDCl_3):



- 7.40 (s, 0.5H, H2)
- 7.12 (s, 1H, H8)
- 6.71 (d, 0.5H, $J=5.6$ Hz, H6)
- 6.35 (d, 0.5H, $J=5.6$ Hz, H5)
- 6.33 [s, 0.5 H, H5(D6)]
- 5.40 (t, 1H, $J=6.8\pm 0.3$ Hz, H10)
- 4.52 (dd, 1H, $J=2.2, 7.6$ Hz, H12)
- 4.33 (d, 1H, $J=11.2$ Hz, H13)
- 4.10 (d, 1H, $J=11.2$ Hz, H13)
- 3.72 (s, 3H, CO_2CH_3)
- 3.22 (AB, 2H, $J=14.4$ Hz, SCH_2)
- 2.50 (ddd, 1H, $J=2.2, 6.5, 8.6$ Hz, H11)
- 2.32-2.42 (m, 1H, H11)
- 1.22 (s, 9H, $\text{O}_2\text{Ct-Bu}$)
- 0.92 (s, 9H, $\text{Si}(\text{CH}_3)_3$)
- 0.17 (s, 3H, SiCH_3)
- 0.14 (s, 3H, SiCH_3)

FTIR (neat film, cm^{-1}): 3485(b), 1732(m), 1716(s), 1532(m)

TLC (1:2 EtOAc:Hexanes): $R_f = 0.37$

HR FAB MS

(matrix: Nitrobenzyl alcohol): calcd: 523.2519 $[\text{MH}]^+$
found: 523.2560



J-1087-1716

J-1087-m17

Proton NMR spectra recorded on a JEOL JNM-GX400 instrument
 Solvent: Acetic acid-d₄. Reference: Acetic acid at 2.04 ppm. Digital resolution: 0.24 Hz

Proton NMR data NCS Chromophore(1)					Proton NMR data of Nitrosoamine(2)				
	δ	m	# of H	<i>J</i> (Hz)		δ	m	# of H	<i>J</i> (Hz)
4''	8.09	d	1H	9.2	4''	8.06	d	1H	8.8
8''	7.89	d	1H	2.2	8''	7.86	d	1H	2.2
3''	7.03	d	1H	9.2	3''	7.02	d	1H	8.8
6''	6.90	d	1H	1.2	6''	6.87	d	1H	1.5
12	6.79	s	1H		12	6.73	s	1H	
11	6.36	s	1H		11	5.92	s	1H	
						[5.96	s	1H] [†]	
8	5.78	s	1H		8	5.72	s	1H	
						[5.71	s	1H] [†]	
1'	5.77	s	1H		1'	5.58	d	1H	3.6
						[5.36	d	1H	3.7] [†]
10	5.08	s	1H		10	4.98	s	1H	
13	4.90	dd	1H	5.5, 8.5	13	4.89	dd	1H	5.5, 8.5
14a	4.72	t	1H	8.5	14a	4.71	t	1H	8.5
14b	4.53	dd	1H	5.5, 8.5	14b	4.54	dd	1H	5.5, 8.5
3'	4.40	dd	1H	3.2, 10.5	3'	4.60	dd	1H	3.8, 11.3
						[4.49	dd	1H	3.8, 11.3] [†]
5'	4.11	q	1H	6.6	5'	4.25	q	1H	6.6
						[4.19	q	1H	6.6] [†]
5	4.08	s	1H		5	4.08	s	1H	
4'	3.93	s	1H		4'	4.06	s	1H	
OCH ₃	3.83	s	3H		OCH ₃	3.81	s	3H	
2'	3.78	dd	1H	3.2, 10.6	2'	5.11	dd	1H	3.7, 11.3
						[5.23	dd	1H	3.7, 11.3] [†]
NCH ₃	3.03	s	3H		NCH ₃	3.24	s	3H	
						[3.96	s	3H] [†]	
5'-CH ₃	1.27	d	3H	6.6	5'-CH ₃	1.33	d	3H	6.6

[†]Minor Rotamer (presumed *syn*)

J-1087-m18

^1H NMR spectra (400 MHz, $\text{CD}_3\text{CO}_2\text{D}$) of neocarcinostatin chromophore (1, bottom) and neocarcinostatin chromophore nitrosamine (9, top). Smaller numerals refer to the *Z*-rotamer, *m* to methylene chloride.

